

LABEL

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

Assurant 200 Insecticide

ACTIVE CONSTITUENT: 200 g/L FIPRONIL

GROUP	2B	INSECTICIDE
--------------	-----------	--------------------

For the control of ants, mole cricket and Argentine stem weevil in commercial turf and the control of various insect pests in asparagus, bananas, brassicas, cotton, forestry, ginger, wine grapevines, mushrooms, pasture, potatoes, sorghum, sugarcane and swede and as specified in the Directions for Use Table

**IMPORTANT:
READ THE ATTACHED LEAFLET BEFORE USE**

Contents: 1, 2, 5, 10, 20 L



Sherwood Chemicals Australasia Pty Ltd
Level 3, 1060 Hay Street
WEST PERTH 6005
AUSTRALIA
Tel: 08 9219 4683
Fax: 08 9219 4672

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container and preparing the product for use, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow length PVC gloves and goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow length PVC gloves. Wash hands after use. After each days use, wash gloves and goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Telephone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS), which can be obtained from your supplier.

CONDITIONS OF SALE

All conditions and warranties rights and remedies implied by law or arising in contract or tort whether due to the negligence of Sherwood Chemicals or otherwise are hereby expressly excluded so far as the same may legally be done provided however that any rights of the Buyer pursuant to non excludable conditions or warranties of the Trade Practices Act 1974 or any relevant legislation of any State are expressly preserved but the liability of Sherwood Chemicals or any intermediate Seller pursuant thereto shall be limited if so permitted by the said legislation to the replacement of the goods sold or the supply of equivalent goods and all liability for indirect or consequential loss or damage of whatsoever nature is expressly excluded. This product must be used or applied strictly in accordance with the instructions appearing hereon. This product is solely sold for use in Australia and must not be exported without the prior written consent of Sherwood Chemicals.

Product Number:

Batch Number:

DOM:

APVMA Approval Number: 66873/106355

Sherwood Chemicals Australasia Pty Ltd

Level 3, 1060 Hay Street

WEST PERTH 6005

AUSTRALIA

Tel: 08 9219 4683

Fax: 08 9219 4672

LEAFLET

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

ASSURANT 200

Insecticide

ACTIVE CONSTITUENT: 200 g/L FIPRONIL

GROUP	2B	INSECTICIDE
-------	-----------	-------------

For the control of ants, mole cricket and Argentine stem weevil in commercial turf and the control of various insect pests in asparagus, bananas, brassicas, cotton, forestry, ginger, wine grapevines, mushrooms, pasture, potatoes, sorghum, sugarcane and swede and as specified in the Directions for Use Table

IMPORTANT:
Read this booklet before use

Contents: 1, 2, 5, 10, 20 L



Sherwood Chemicals Australasia Pty Ltd
Level 3, 1060 Hay Street
WEST PERTH 6005
AUSTRALIA
Tel: 08 9219 4683
Fax: 08 9219 4672

Website: www.sherwoodchemicals.com.au

APVMA Approval Number: 66873/106355

DIRECTIONS FOR USE

Crop	Pest	State	Rate	WHP	Critical Comments
					Dangerous to bees. Refer to PROTECTION OF LIVESTOCK.
Asparagus	Garden weevil (<i>Phlyctinus callosus</i>)	WA, Vic only	40 mL/100 L applying 500 L spray solution per hectare	1 day (H)	Day time spraying is effective but superior control may be achieved if spray is applied at night. Repeat applications as required, depending on pest pressure. Controlling weevils in asparagus fern may reduce abundance the following season. DO NOT apply more than 6 applications per season, up to 4 applications to spears and after harvest 2 applications.
Bananas	Banana rust thrips (<i>Chaetanapho- thrips signipennis</i>)	Qld, NSW, WA, NT only	Butt application All planting configurations: 150 mL/100 L water (0.75 mL/stool) Band application All planting configurations: 40 mL/100 m ² treated area (See table under 'General Instructions - Application' for calculation of the treated area)	-	Timing Application should ideally be made at least two months prior to bunch emergence to reduce early thrips pressure. Such an application could coincide with an application for banana weevil borer control (see label directions below). Butt application Apply in a coarse spray covering the stem to a height of 30 cm and the soil/trash in a 30 cm radius from the stem base. Apply a total volume of 500 mL solution per stool. Ensure thorough coverage of butt, suckers, trash and exposed soil. Band application Apply in a band along each row. The band width should be such that at least 30 cm of soil/trash is treated on both sides of the butt. Apply with a side delivery boom and offset nozzles directed to spray at least 30 cm of soil on either side of the butt and to a height of 30 cm up the stems. Repeat the application from the opposite side of the row. Half of the spray volume required to treat each row should be applied from each direction of spraying. For double row configurations, treat both rows with each pass, ensuring the ground area between the two rows is also treated. Ensure thorough coverage of butt, suckers, trash and exposed soil. Apply in a minimum water volume of 13 L/100 m ² (trash removed) or 26 L/100 m ² (trash retained). See table under 'General Instructions – Application' for guidance.
	Banana weevil borer (<i>Cosmopolites sordidus</i>)	Qld, NSW, WA, NT only	Butt application All planting configurations 150 mL/100 L water (0.75 mL/stool)	-	Apply by butt application as described above for banana rust thrips. Population assessment Lay baits (cut billets of stem base) flat on soil beside stools and cover with leaf material. Check baits after 3 days to assess pest activity. Monitoring should commence in September when pest activity increases and continue until April. Application method Applications should be made in spring and/or autumn when weevil numbers reach or exceed acceptable threshold levels. Remove any green trash from area to be treated. Avoid application to trash, which is less than 3 weeks old. This use is subject to a Croplife Resistance Management Strategy. Refer to your Sherwood representative for details.

Brassicas (head cabbage, cauliflower, broccoli, Brussels sprout, kohlrabi)					Dangerous to bees. Refer to PROTECTION OF LIVESTOCK.
	Diamondback moth (<i>Plutella xylostella</i>), cabbage white butterfly (<i>Pieris rapae</i>), cabbage cluster caterpillar (<i>Crociodolomia pavonana</i>)	All States	250 mL/ha	7 days (H)	Diamondback moth can rapidly become resistant to insecticides. To preserve the effectiveness of this product, limit the number of applications to no more than 4 per year, preferably applied within an 8 week period. Use spray volume of between 400 and 1000 L/ha according to crop size. Use a non-ionic wetting agent at the rate specified by the manufacturer for use in horticultural crops. Ensure that the rate of wetting agent used results in efficient spray coverage of the leaf surface. This use is subject to a Croplife Resistance Management Strategy. Refer to your Sherwood representative for details.
Cotton	Cotton thrips (<i>Thrips tabaci</i>) Green mirid (<i>Creontiades dilutus</i>)	Qld, NSW, WA only	62.5 to 125 mL/ha	4 weeks (H)	Apply at the first sign of the pest. Assurant 200 will take 3-4 days to reach full effectiveness. Apply spray to achieve thorough coverage of foliage when pest first appears and repeat as required. Use the higher rate in situations of high thrips pressure. Use the higher rate under sustained heavy green mirid pressure. Use higher rate in situations of high green vegetable bug pressure. The product is compatible with early season IPM with the lower rate having less impact on beneficials.
Forestry Plantations including Eucalyptus, Pinus and Corymbia spp.	Australian plague locust (<i>Chortoicetes terminifera</i>) Spur-throated locust (<i>Austracris guttulosa</i>) Migratory locust (<i>Locusta migratoria</i>) Wingless grasshopper (<i>Phaulacridium vittatum</i>) Small plague grasshopper (<i>Austroicetes cruciata</i>)	All states	62.5 to 125 mL/ha	-	Apply in plantations situations up to a maximum of two years of age. Apply diluted with water to a minimum of 20 L/ha by air or 50 L/ha by ground rig, directly onto locusts. Ensure thorough coverage of foliage. Residual control of these pests provided by Assurant 200 will vary with conditions. Rainfall will significantly reduce residual control. Residual control will also be reduced when applied directly to bare earth. Where inaccessibility prevents direct spraying of locusts apply as a barrier treatment (minimum 25 m wide) ahead of advancing hopper bands. DO NOT retreat for 14 days following application. DO NOT apply Assurant 200 to wet foliage. Assurant 200 is rainfast after drying on foliage (1 hour). Re-spray only if rain falls before spray is dry on crop. Mortality will increase to a maximum over a period of 3-15 days after spraying. Speed of kill varies with locust species, temperature and age of adults. Fully mature, adult spur-throated locusts may show symptoms of debilitation 4-48 hours after spraying but in cool weather may take up to 14 days to die. Feeding ceases when debilitation symptoms appear. See also General Instructions .
Ginger	Symphylids	Qld only	Pre-plant application 250 mL to 500 mL/ha Dipping 1 mL/200 L + pre-planting application	-	Pre-plant application Apply as a spray to soil and incorporate to a depth of 200mm prior to planting. Use the higher rate in heavier soils and/or under high pest pressure. Dipping Dip seed pieces in aqueous solution prior to planting in beds pre-treated with Assurant 200. Dip solution is achieved by mixing 1 mL of Assurant 200 in 200 L of water.

					Dangerous to bees. Refer to PROTECTION OF LIVESTOCK.
Wine grapevines	Fig longicorn (<i>Acalolepta vastator</i>)	All States	100 mL/100 L	-	Apply as a single spray to dormant vines following pruning and prior to budburst. Apply only as a high volume spray using hand held equipment. Thorough coverage of vine trunks and cordons is essential for effective control. Refer to Application Wine grapevines.
Mushrooms	Mushroom flies (<i>Sciarids, Phorids and Cecids</i>)	All States	16 mL/300 L bale of peat moss	14 days (H)	Prepare solution by mixing Assurant 200 with a small volume of water. Apply mixture to peat moss during preparation of casing. Ensure thorough mixing with peat moss.
Pasture, Sorghum	Australian plague locust (<i>Chortoicetes terminifera</i>), Spur-throated locust (<i>Austracris guttulosa</i>), Migratory locust (<i>Locusta migratoria</i>) Wingless grasshopper (<i>Phaulacridium vittatum</i>) Small plague grasshopper (<i>Austroicetes cruciata</i>)	All States	6.25 mL/ha	14 days (H, G)	Apply diluted with water to a minimum of 20 L/ha by air or 50 L/ha by ground rig, directly onto locusts. Ensure thorough coverage of foliage. Residual control of these pests provided by Assurant 200 will vary with conditions. Rainfall will significantly reduce residual control. Residual control will also be reduced when applied directly to bare earth. Where inaccessibility prevents direct spraying of locusts apply as a barrier treatment (minimum 25 m wide) ahead of advancing hopper bands. DO NOT retreat for 14 days following application. DO NOT apply Assurant 200 to wet foliage. Assurant 200 is rainfast after drying on foliage (1 hour). Re-spray only if rain falls before spray is dry on crop. Mortality will increase to a maximum over a period of 3-15 days after spraying. Speed of kill varies with locust species, temperature and age of adults. Fully mature, adult spur-throated locusts may show symptoms of debilitation 4-48 hours after spraying but in cool weather may take up to 14 days to die. Feeding ceases when debilitation symptoms appear. See also General Instructions.
Potatoes, Sweet potatoes	Wireworm (various), mole cricket (various)	All states	250 mL/ha	-	Apply as a broadcast spray to the surface of the soil and incorporate to a depth of 15 cm prior to planting.
	Whitefringed weevil (<i>Naupactus leucoloma</i>)		500 mL/ha		

Sugarcane					Dangerous to bees. Refer to PROTECTION OF LIVESTOCK.
	Sugarcane weevil borer (<i>Rhabdoscelus obscurus</i>)	Qld, NSW, WA, NT only	2 to 5.7 mL/100 m row	12 weeks (H, G)	Apply during the summer months of December to February when the crop has produced the first millable internode of cane. Use hollow cone nozzles as a directed spray to cover the base of the sugarcane stools and up the stalk to a height of 40 cm. Treat both sides of the stools ensuring coverage of all stalks, soil and trash in an area to 10 cm either side of the stools. Use a non-ionic wetting agent at the rate specified by the manufacturer. Ensure that the rate of wetting agent used results in efficient spray coverage of the stalk, soil and trash surface. Apply in a minimum water volume of 250 L/ha (approx. 3.8 L/100 m row). Use the higher rate when pest pressure is heavy.
	Sugarcane wireworm (various)		Single row plantings: 1.1 mL/100 m single row length Double row plantings: 1.8 mL/100 m double row length 250 mL/ha	-	Apply in the planting furrow over the top of the plant pieces (setts), in sufficient water to ensure coverage of the plant pieces and the surrounding soil.
Swede Turnips	Diamondback moth (<i>Plutella xylostella</i>)	All states	250 mL/ha	7 days (H)	Use according to the CropLife Resistance Management Strategy for diamondback moth control. The use is limited to 4 applications per year, preferably applied within an 8 week period.

DIRECTIONS FOR USE – Turf			
Pest	Situation	Rate	Critical Comments
Ant pests of turf, including but not limited to: Funnel ant (<i>Aphaenogaster</i> spp.), greenhead ants (<i>Rhytidoponera</i> spp.) Argentine stem weevil (<i>Listrionotus bonariensis</i>) Mole cricket (<i>Gryllotalpa</i> spp.)	Turf (eg. lawns, commercial turf farms, parks, recreational areas, bowling greens, golf courses, sports fields)	300 mL/ha (3 mL/100 m ²)	Mix Assurant 200 in water and apply a minimum spray volume of 300 L/ha (3 L/100 m ²) evenly over the area to be treated using spray equipment that delivers a coarse droplet size at the first sign of pest activity. To ensure optimum control, irrigate the treated area with up to 4-6 mm of water soon after application.
		375 mL/ha	Apply to areas where ants are active. Where possible spray directly into the nests. Initial applications should be broadcast over affected areas as ant activity increases in early spring. If required, a repeat application may be made during summer.
		150 mL/ha	
			In domestic turf situations, use is only by authorised lawn care specialists, professional pest managers or persons with relevant chemical handling certification.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

WITHHOLDING PERIODS:

(H) = Harvest, (G) = Grazing

ASPARAGUS:	DO NOT HARVEST FOR 1 DAY AFTER APPLICATION
BANANAS:	NOT REQUIRED WHEN USED AS DIRECTED
BRASSICAS:	DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION
COTTON:	DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION DO NOT GRAZE OR CUT FOR STOCK FOOD
WINE GRAPEVINES:	NOT REQUIRED WHEN USED AS DIRECTED (H) DO NOT FEED TRASH OR BY-PRODUCTS RESULTING FROM TREATED GRAPEVINES TO LIVESTOCK (G)
GINGER:	NOT REQUIRED WHEN USED AS DIRECTED
MUSHROOMS:	DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION
PASTURE:	DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION
POTATOES:	NOT REQUIRED WHEN USED AS DIRECTED DO NOT GRAZE OR CUT FOR STOCK FOOD ANY PART OF FAILED CROP (INCLUDING TUBERS)
SORGHUM:	DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION
SUGARCANE:	DO NOT HARVEST FOR 12 WEEKS AFTER APPLICATION. DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 12 WEEKS AFTER APPLICATION
SWEDE AND TURNIPS:	DO NOT HARVEST FOR 7 DAYS FOLLOWING APPLICATION DO NOT ALLOW LIVESTOCK TO GRAZE TREATED CROP
POTATOES:	NOT REQUIRED WHEN USED AS DIRECTED DO NOT GRAZE OR CUT FOR STOCK FOOD ANY PART OF FAILED CROP (INCLUDING TUBERS)
TURF:	DO NOT GRAZE TREATED AREAS OR FEED GRASS CLIPPINGS FROM TREATED AREAS TO ANIMALS INCLUDING POULTRY

LIVESTOCK WITHHOLDING PERIOD

WITHHOLD STOCK FROM SLAUGHTER FOR 21 DAYS AFTER APPLICATION, WHERE STOCK WERE PRESENT IN CROP OR PASTURE AT TIME OF APPLICATION

GENERAL INSTRUCTIONS**INSECTICIDE RESISTANCE WARNING**

GROUP	2B	INSECTICIDE
--------------	-----------	--------------------

For insecticide resistance management Assurant 200 is a Group 2B insecticide. Some naturally occurring insect biotypes resistant to Assurant 200 and other Group 2B insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Assurant 200 or other Group 2B insecticides are used repeatedly. The effectiveness of Assurant 200 on resistant individuals could be significantly reduced. Since the occurrence of resistant individuals is difficult to detect prior to use, Sherwood Chemicals Australasia Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant insects. Assurant 200 may be subject to specific resistance management strategies. For further information contact your local supplier, Sherwood representative or local agricultural department agronomist.

RESIDUES MANAGEMENT IN EXPORT PRODUCE**Crops**

Growers should note that MRLs or import tolerances do not exist in all markets for produce treated with Assurant 200. If you are growing produce for export, please check with Sherwood Chemicals Australasia Pty Ltd for the latest information on MRLs and import tolerances BEFORE using Assurant 200.

Livestock

Livestock may be exposed to Assurant 200 residues in the feed by grazing treated pasture and/or sorghum forage and fodder. Observance of the 14 day grazing withholding period permits compliance with Australian MRLs for Assurant 200 in meat, offal and milk. To meet more stringent export residues requirements, Meat and Livestock Australia recommends Export Slaughter Intervals (ESIs) and Export Grazing Intervals (EGIs) for Assurant 200. When livestock grown for export are grazed on pasture and/or sorghum forage and fodder treated with Assurant 200 the user must obtain details of the recommended export intervals from Meat and Livestock Australia and must follow those recommendations.

MIXING

Asparagus, Bananas, Brassicas, Cotton, Ginger, Wine Grapevines, Pasture, Potatoes, Sweet Potatoes, Sorghum, Sugarcane

Slowly add the required amount of product to water in the spray tank while stirring or agitating. Agitate while spraying.

Mushrooms

Add required amount of product to a small quantity of water, ensuring thorough mixing.

APPLICATION**Bananas**

Ensure thorough coverage of butts, suckers and surrounding trash and exposed soil.

Bananas (Band Spray)

Example calculations of the quantity of Assurant 200 and the minimum water volume required to treat a 100 m row length of bananas for various band widths:

Band width to be Treated #	Spray area per 100 m row	Total quantity of Assurant 200 required per 100 m row *	Minimum recommended water volume per 100 m (trash removed) *	Minimum recommended water volume per 100 m (trash retained) *
1.5 m	150 m ²	60 mL	20 L	40 L
2.0 m	200 m ²	80 mL	27 L	54 L
2.5 m	250 m ²	100 mL	33 L	66 L
3.0 m	300 m ²	120 mL	40 L	80 L

Band width = butt diameter plus 30 cm on either side of the butt

* Rows should be treated from both sides. The quantities stated are the total amounts to be applied, i.e. half of the stated quantity should be applied from each direction of spraying.

Brassicas

Ensure thorough coverage of foliage and heads.

Cotton

For ground application use a prepared spray volume of 35 – 75 L/ha depending on the size of the crop. For aerial application see 'Aerial application' instructions below.

Ginger

Apply as a spray to soil and incorporate to a depth of 200 mm prior to planting.

Wine Grapevines

Assurant 200 should be applied by hand held equipment as a high volume directed spray of approximately 500 mL of solution per vine directed to trunks and cordons during dormant period only.

Mushrooms

Apply mixture to peat moss during preparation of casing, ensuring even mixing in peat moss.

Potatoes, Sweet Potatoes

Apply as a broadcast spray to the surface of the soil and incorporate to a depth of 15 cm prior to planting.

Swede (including Turnip and Radish)

Ensure thorough coverage of foliage.

Turf

Apply a minimum spray volume of 300 L/ha (3 L/100 m²) evenly over the area to be treated using spray equipment that delivers a coarse droplet size at the first sign of pest activity.

DO NOT apply with aircraft or through any type of irrigation equipment.

DO NOT apply this product using hand held equipment.

DO NOT apply more than twice a year.

AERIAL APPLICATION

Use spray techniques that minimise off-target spray drift. DO NOT use rotary atomisers. Use application volumes between 20 L and 50 L/ha. Achieve a droplet density of ~60 droplets/cm² on a flat surface on the target. When spraying large droplets (>250 µm), increase the application volume to >40 L/ha to ensure sufficient droplets are produced. Aerial application is not recommended for Brassica, potato crops and turf uses.

PRECAUTIONS

Re-entry period

DO NOT allow entry into treated areas until spray has dried. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). Clothing must be laundered after each day's use. Human flaggers, if used in aerial spraying operations, must be protected by enclosed cabs.

RE-ENTRY – TURF ONLY

Public

DO NOT allow public access until spray has dried.

Workers

DO NOT allow entry into treated areas for nine days for low exposure activities such as irrigation and mowing, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

DO NOT allow entry into treated areas for 37 days for high exposure activities such as hand weeding and transplanting, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply in weather conditions or from spraying equipment that may cause spray to drift onto non-target plants/crops, cropping lands or pastures.

PROTECTION OF LIVESTOCK

Dangerous to bees. DO NOT apply where bees from managed hives are known to be foraging, and crops, weeds or cover crops are in flower at the time of spraying, or are expected to flower within 28 days (7 days for pastures and sorghum).

Before spraying, notify beekeepers to move hives to a safe location with an untreated source of nectar, if there is potential for managed bees to be affected by the spray or spray drift. If an area has been sprayed inadvertently, in which the crop, weeds or cover crop were in flower or subsequently came into flower, notify beekeepers in order to keep managed bees out of the area for at least 28 days

(7 days for pastures and sorghum) from the time of spraying. Where the owner of managed hives in the vicinity of a crop to be sprayed is not known, contact your State Department of Primary Industries/Agriculture, citing the registration number, for assistance in contacting the owner.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT spray across open bodies of water. Highly toxic to fish and aquatic organisms. This product will kill susceptible non-target invertebrates, including beneficial species, if they are exposed to drift. DO NOT apply aerially to brassicas and potatoes. A spray drift minimisation strategy should be employed at all times when aerially applying sprays to, or near, sensitive areas. The strategy envisaged is exemplified by the cotton industry's Best Management Practice Manual.

STORAGE AND DISPOSAL

2.5 L, 5 L, 10 L, 20 L

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods indirect sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container and preparing the product for use, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow length PVC gloves and goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow length PVC gloves. Wash hands after use. After each days use, wash gloves and goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Telephone Australia 13 11 26.

MSDS

For further information refer to the Material Safety Data Sheet (MSDS), which can be obtained from your supplier.

CONDITIONS OF SALE

All conditions and warranties rights and remedies implied by law or arising in contract or tort whether due to the negligence of Sherwood Chemicals or otherwise are hereby expressly excluded so far as the same may legally be done provided however that any rights of the Buyer pursuant to non excludable conditions or warranties of the Trade Practices Act 1974 or any relevant legislation of any State are expressly preserved but the liability of Sherwood Chemicals or any intermediate Seller pursuant thereto shall be limited if so permitted by the said legislation to the replacement of the goods sold or the supply of equivalent goods and all liability for indirect or consequential loss or damage of whatsoever nature is expressly excluded. This product must be used or applied strictly in accordance with the instructions appearing hereon. This product is solely sold for use in Australia and must not be exported without the prior written consent of Sherwood Chemicals.

APVMA Approval Number: 66873/106355